arXiv:1209.5986v1 [math.PR] 26 Sep 2012

Signal reconstruction from the magnitude of subspace components

C. Bachoc[a], M. Ehler[b]

[a] Univ. Bordeaux, IMB, UMR 5251, F-33400 Talence, France
[b] Helmholtz Zentrum München, Institute of Biomathematics and Biometry, Ingolstädter Landstrasse 1, 85764 Neuherberg, Germany

---

Abstract

We consider signal reconstruction from the norms of subspace components. If the weighted linear subspaces form a tight p-fusion frame and a cubature formula of strength 4, then even in case that p of the subspace norms are erased, we find a finite list of potential signals, one of which is the correct one. Moreover, we present a computationally feasible algorithm to determine this list. Alternatively and to reduce the number of required measurements to a linear scale in the ambient dimension, we use semidefinite programming and random subspaces.

---

1. Introduction

Frames have become a powerful tool in signal processing that can offer more flexibility than orthogonal bases, cf. [18, 42, 41] and [26, 27, 28, 30]. Many signal processing problems in engineering such as X-ray crystallography and diffraction imaging require signal reconstruction from the magnitude of its frame coefficients, also known under the terms *phase retrieval* and *reconstruction without the phase*, cf. [7, 11] and references therein. By making additional assumptions on the underlying frame, exact solutions are presented in [6, 7], see [45, 46] for relations to quantum measurements. Recently, reconstruction from frame coefficients without phase has been numerically addressed via semidefinit programming, see [11, 13].

Frame coefficients can be thought of as projections onto 1-dimensional subspaces. In image reconstruction from averaged diffraction patterns by means of incoherent addition of k wavefields [32], the original signal must be recovered from the norms of its k-dimensional subspace components. Notably, the latter is a common problem in crystal twinning [24]. Here, we pose the following questions: Can we reconstruct the original signal from the norms of its k-dimensional

---

Email addresses: `Christine.Bachoc@math.u-bordeaux1.fr` (C. Bachoc), `martin.ehler@helmholtz-muenchen.de` (M. Ehler)

subspace components by means of a closed formula? Also, can we reconstruct even if a certain number of norms of subspace components are erased?

In this note we shall provide an affirmative answer in the following sense: Given k-dimensional linear subspaces $\{V_j\}_{j=1}^n$ in $\mathbb{R}^d$, we aim to reconstruct the signal $x \in S^{d-1}$ from any subset of $\{\|P_{V_j}(x)\|\}_{j=1}^n$ that has cardinality at least $n-p$. Here, P_{V_j} denotes the orthogonal projection onto V_j. Under suitable conditions on the subspaces, we are able to compute a finite list of candidate signals, one of which is the correct one.

Finding a set of candidate signals is known as list decoding and was introduced in [31]. We determine this list in a two-step procedure. Without loss of generality, let us suppose that the first p norms were erased. If there are positive weights $\{\omega_j\}_{j=1}^n$ such that $\{(V_j, \omega_j)\}_{j=1}^n$ forms a tight p-fusion frame as recently introduced in [5], then we are able to reconstruct the erased norms $\{\|P_{V_j}(x)\|\}_{j=1}^p$ at least up to permutations. Here, we need to verify that a certain system of algebraic equations has only finitely many solutions. Notice that our input are not the subspace components but their norms, as opposed to signal reconstruction under the erasures discussed in [10, 14, 40, 43].

In the second step, we assume that $\{(V_j, \omega_j)\}_{j=1}^n$ yields a cubature formula of strength 4, see [5], enabling us to reconstruct the orthogonal projection onto $x\mathbb{R}$ from knowledge of $\{\|P_{V_j}(x)\|\}_{j=1}^n$. This extends the 1-dimensional results in [6] to k-dimensional projections addressed in the present paper. Note that the authors in [6] require cubature formulas for the projective space whose weights are $\omega_j = 1/n$, i.e., so-called projective designs. In practice, however, the choice of subspaces may underlie some restrictions that prevent them from being a design. Nevertheless, given subspaces sufficiently dense in the Grassmann space, nonnegative weights can be computed so that we obtain a cubature of a certain strength depending on the density to begin with, cf. [33, 34, 36]. Thus, our results in Step 2 are a significant improvement for 1-dimensional projections already.

The cardinality of a cubature formula of strength 4 scales at least quadratically with the ambient dimension d. For $k = 1$, it is known that semidefinite programming yields signal recovery with high probability when the 1-dimensional subspaces are chosen at random [13]. The cardinality of the subspaces can then scale linearly in the ambient dimension up to a logarithmic factor. This factor has recently been removed in [12]. Here, we shall extend such results to general k-dimensional subspaces chosen at random. Our proof is guided by the approach in [13] but some steps are more involved and need to be taken care of with different tools in this more general setting. For instance, the case $k = 1$ relies on matrices whose entries are i.i.d. Gaussian modeling the measurements. For $k > 1$, we must deal with measurement matrices involving orthogonal projectors that clearly have dependent entries. Hence, the extension from $k = 1$ to $k > 1$ is not obvious and requires special care. The stability under noise, on the other hand, follows from the lines in [13]. We also verify numerically that semidefinite programming enables us to recover a signal from the norms of its random k-dimensional subspace components.

The outline is as follows: In Section 2, we recall fusion frames and state the problem. In Section 3, we introduce tight p-fusion frames and cubature formulas. We present our reconstruction algorithm in Section 4. Semidefinite programming for signal reconstruction is addressed in Section 5, and its stability under noise is discussed in Section 6. In Section 7, the numerical experiments using semidefinite programming are presented. Conclusions are given in Section 8.

2. Fusion frames and the problem of reconstruction without phase

Let $\mathcal{G}_{k,d} = \mathcal{G}_{k,d}(\mathbb{R})$ denote the *real Grassmann space*, i.e., the k-dimensional subspaces of $\mathbb{R}^d$. Each $V \in \mathcal{G}_{k,d}$ can be identified with the orthogonal projection onto V, denoted by P_V. Let $\{V_j\}_{j=1}^n \subset \mathcal{G}_{k,d}$ and let $\{\omega_j\}_{j=1}^n$ be a collection of positive weights. Then $\{(V_j, \omega_j)\}_{j=1}^n$ is called a *fusion frame* if there are positive constants A and B such that

$$A\|x\|^2 \leq \sum_{j=1}^n \omega_j \|P_{V_j}(x)\|^2 \leq B\|x\|^2, \text{ for all } x \in \mathbb{R}^d, \tag{1}$$

cf. [15]. The condition (1) is equivalent to

$$A \leq \sum_{j=1}^n \omega_j \langle P_x, P_{V_j} \rangle \leq B, \text{ for all } x \in S^{d-1}, \tag{2}$$

where P_x is short for $P_{x\mathbb{R}^d}$ and $\langle P_x, P_{V_j} \rangle := \operatorname{trace}(P_x P_{V_j})$ is the standard inner product between self-adjoint operators. If $A = B$, then $\{(V_j, \omega_j)\}_{j=1}^n$ is called a *tight fusion frame*, and any signal $x \in S^{d-1}$ can be reconstructed from its subspace components by the simple formula

$$x = \frac{1}{A} \sum_{j=1}^n \omega_j P_{V_j}(x). \tag{3}$$

If, however, instead of $\{P_{V_j}(x)\}_{j=1}^n$ we only observe the norms $\{\|P_{V_j}(x)\|\}_{j=1}^n$ and, worse, we even lose some of these norms, can we still reconstruct x? In the present paper, we find conditions on $\{(V_j, \omega_j)\}_{j=1}^n$ together with a computationally feasible algorithm such that we can determine a finite list of candidate signals, one of which is the correct one.

3. Tight p-fusion frames and cubature formulas

3.1. Tight p-fusion frames

Let $\{V_j\}_{j=1}^n \subset \mathcal{G}_{k,d}$ and let $\{\omega_j\}_{j=1}^n$ be a collection of positive weights and p a positive integer. Then $\{(V_j, \omega_j)\}_{j=1}^n$ is called a *p-fusion frame* in [5] if there exist positive constants A and B such that

$$A\|x\|^{2p} \leq \sum_{j=1}^n \omega_j \|P_{V_j}(x)\|^{2p} \leq B\|x\|^{2p}, \text{ for all } x \in \mathbb{R}^d. \tag{4}$$

If $A = B$, then $\{(V_j, \omega_j)\}_{j=1}^n$ is called a *tight p-fusion frame*. As with (1) and (2), the condition (4) is equivalent to

$$A \leq \sum_{j=1}^n \omega_j \langle P_x, P_{V_j}\rangle^p \leq B, \text{ for all } x \in S^{d-1}. \tag{5}$$

If $\{(V_j, \omega_j)\}_{j=1}^n$ is a tight p-fusion frame, then it is also a tight ℓ-fusion frame for all integers $1 \leq \ell \leq p$, and the tight ℓ-fusion frame bounds are

$$A_\ell = \frac{(k/2)_\ell}{(d/2)_\ell} \sum_{j=1}^n \omega_j, \tag{6}$$

where we used $(a)_\ell = a(a+1)\cdots(a+\ell-1)$, cf. [5]. We also refer to [5] for constructions and general existence results.

3.2. Cubature formulas

The real orthogonal group $O(\mathbb{R}^d)$ acts transitively on $\mathcal{G}_{k,d}$, and the Haar measure on $O(\mathbb{R}^d)$ induces a probability measure σ_k on $\mathcal{G}_{k,d}$. Let $L^2(\mathcal{G}_{k,d})$ denote the complex valued functions on $\mathcal{G}_{k,d}$, whose squared module is integrable with respect to σ_k. The complex irreducible representations of $O(\mathbb{R}^d)$ are associated to partitions $\mu = (\mu_1, \ldots, \mu_d)$, $\mu_1 \geq \ldots \geq \mu_d \geq 0$, denoted by V_d^μ, cf. [37]. Let $l(\mu)$ be the number of nonzero entries in μ so that

$$L^2(\mathcal{G}_{k,d}) = \bigoplus_{l(\mu)\leq k} H_{k,d}^{2\mu}, \qquad \text{where } H_{k,d}^{2\mu} \simeq V_d^{2\mu}, \tag{7}$$

see [37]. The space of polynomial functions on $\mathcal{G}_{k,d}$ of degree bounded by $2p$ is

$$\mathrm{Pol}_{\leq 2p}(\mathcal{G}_{k,d}) := \bigoplus_{l(\mu)\leq k,\ |\mu|\leq p} H_{k,d}^{2\mu}, \tag{8}$$

and we additionally define the subspace

$$\mathrm{Pol}^1_{\leq 2p}(\mathcal{G}_{k,d}) := \bigoplus_{l(\mu)\leq 1,\ |\mu|\leq p} H_{k,d}^{2\mu}. \tag{9}$$

Let $\{V_j\}_{j=1}^n \subset \mathcal{G}_{k,d}$ and $\{\omega_j\}_{j=1}^n$ be a collection of positive weights normalized such that $\sum_{j=1}^n \omega_j = 1$. Then $\{(V_j, \omega_j)\}_{j=1}^n$ is called a *cubature of strength* $2p$ *for* $\mathcal{G}_{k,d}$ if

$$\int_{\mathcal{G}_{k,d}} f(V) d\sigma_k(V) = \sum_{j=1}^n \omega_j f(V_j) \qquad \text{for all } f \in \mathrm{Pol}_{\leq 2p}(\mathcal{G}_{k,d}). \tag{10}$$

Grassmannian designs, i.e., cubatures with constant weights, have been studied in [1, 2, 3, 4]. For existence results on cubatures and the relations between p

and n, we refer to [22]. It was verified in [5] that $\{(V_j,\omega_j)\}_{j=1}^n$ is a tight p-fusion frame if and only if

$$\int_{\mathcal{G}_{k,d}} f(V)d\sigma_k(V) = \sum_{j=1}^n \omega_j f(V_j) \quad \text{for all } f \in \mathrm{Pol}^1_{\leq 2p}(\mathcal{G}_{k,d}).$$

Thus, any cubature of strength $2p$ is a tight p-fusion frame. The converse implication does not hold in general except for p or k equals 1.

4. Algorithm for signal reconstruction from magnitudes of incomplete subspace components

Let $\{(V_j,\omega_j)\}_{j=1}^n$ be a tight p-fusion frame with $\sum_{j=1}^n \omega_j = 1$ and let $x \in S^{d-1}$. We aim to reconstruct $\pm x$ from only $n-p$ elements of the norms $\{\|P_{V_j}(x)\|^2\}_{j=1}^n$. Without loss of generality, we assume that the first p norms have been erased, so we want to recover $\pm x$ from the knowledge of $\{\|P_{V_j}(x)\|\}_{j=p+1}^n$.

In a first step, we attempt to compute the missing values $t_j := \|P_{V_j}(x)\|^2$, $1 \leq j \leq p$. The second step is dedicated to reconstructing $\pm x$ from $\{\|P_{V_j}(x)\|^2\}_{j=1}^n$.

4.1. Step 1: reconstruction of the erased norms

The tight p-fusion frame $\{(V_j,\omega_j)\}_{j=1}^n$ is also a tight ℓ-fusion frame for $1 \leq \ell \leq p$, so that (6) yields

$$\sum_{j=1}^n \omega_j \|P_{V_j}(x)\|^{2\ell} = A_\ell = \frac{(k/2)_\ell}{(d/2)_\ell}, \quad 1 \leq \ell \leq p.$$

Therefore, $(t_1,\ldots,t_p)$ is a solution of the algebraic system of equations

$$\sum_{j=1}^p \omega_j T_j^\ell = \frac{(k/2)_\ell}{(d/2)_\ell} - \sum_{j=p+1}^n \omega_j \|P_{V_j}(x)\|^{2\ell}, \quad 1 \leq \ell \leq p, \tag{S}$$

in the unknowns $(T_1,\ldots,T_p)$. In the special case of equal weights, (S) gives the values of the symmetric powers $\sum_{j=1}^p t_j^\ell$, for $\ell = 0,\ldots,p$, which, as polynomial expressions, generate the ring of symmetric polynomials up to degree p. Vieta's formula yields

$$\prod_{i=1}^p (T-t_i) = \sum_{j=0}^p (-1)^j e_j T^{p-j}, \quad \text{where } e_0 = 1 \text{ and } e_j = \sum_{1\leq i_1<\ldots<i_j\leq p} t_{i_1}\cdots t_{i_p},$$

and Newton's identity leads to

$$e_j = \frac{1}{j}\sum_{\ell=1}^j (-1)^{\ell-1} e_{j-\ell} \sum_{j=1}^p t_j^\ell, \quad \text{for } j = 1,\ldots,p.$$

Therefore, we can compute the coefficients of $\prod_{i=1}^{p}(T - t_i)$ as a polynomial in T and solve for its roots; we see that $(t_1, \ldots, t_p)$ is determined up to a permutation so that we obtain at most $p!$ distinct solutions to (S).

If the weights are not equal, one can still show that (S) has at most $p!$ solutions. The issue is to verify that the associated affine variety is zero-dimensional. Results from intersection theory and the refined Bézout theorem, cf. [35, 47] and [9], then imply that the variety's cardinality is at most the product of the degrees of the p polynomials, i.e., there are at most $p!$ solutions.

Proposition 4.1. *Let $\{b_\ell\}_{\ell=1}^{p}$ be complex numbers and define*

$$f_\ell(T) = \sum_{j=1}^{p} \omega_j T_j^\ell - b_\ell, \quad \ell = 1, \ldots, p.$$

If $\{\omega_j\}_{j=1}^{p}$ are positive numbers, then the affine variety $\mathcal{V} := \{T \in \mathbb{C}^p : f_1(T) = 0, \ldots, f_p(T) = 0\}$ is zero-dimensional.

Proof. We proceed by induction on p. The assertion is certainly true for $p = 1$. We observe that the Jacobian determinant satisfies

$$\det\left(\frac{\partial(f_1, \ldots, f_p)}{\partial(T_1, \ldots, T_p)}\right) = \omega_1 \cdots \omega_p \cdot p! \cdot \prod_{1 \le i < j \le p} (T_j - T_i),$$

where we used the well-known formula for the Vandermonde determinant. The Jacobian determinant is nonzero for $T \notin \bigcup_{i<j} \Delta_{i,j}$. Therefore, every $T \in \mathcal{V} \setminus \bigcup_{i<j} \Delta_{i,j}$ is a nonsingular point of $\mathcal{V}$, and the dimension of $\mathcal{V}$ at T is $p - p = 0$, cf. [20, Theorem 9.9] and [9, Lemma 11.5.1]. It remains to consider the intersection of $\mathcal{V}$ with the diagonals $\Delta_{i,j} := \{T \in \mathbb{C}^d : T_i = T_j\}$, $i < j$. To fix ideas, let us consider the case $i = 1$, $j = 2$. The intersection $\mathcal{V} \cap \Delta_{1,2}$ is given by the system of equations

$$(\omega_1 + \omega_2) T_2^\ell + \sum_{j=3}^{p} \omega_j T_j^\ell = b_\ell, \quad \ell = 1, \ldots, p.$$

Because $\omega_1 + \omega_2 > 0$, by induction the first $p - 1$ of these equations have only finitely many solutions. Thus, $\mathcal{V} \cap \Delta_{1,2}$ is finite, too. $\square$

We have proved that the system of algebraic equations (S) has at most $p!$ complex solutions. In order to compute these solutions, standard algorithmic methods can be applied [19, 20]. The construction of a Gröbner basis of the ideal $\mathcal{I}$ generated by the p equations allows to compute the algebraic operations in the quotient ring $\mathbb{R}[T_1, \ldots, T_p]/\mathcal{I}$, which is finite dimensional and of dimension at most $p!$. The computation of the solutions then boils down to linear algebra in this space.

4.2. Step 2: reconstruction from the magnitude of subspace components

In the second step, we try to compute P_x from each of the possible candidates for $\{\|P_{V_j}(x)\|^2\}_{j=1}^n$. To that end, we establish the following result that generalizes the case $k = 1$ treated in [6]. We point out that we allow for cubatures as opposed to projective designs in [6] that require the cubature weights to be constant:

Proposition 4.2. *Let $\{(V_j, \omega_j)\}_{j=1}^n$ be a cubature of strength 4 for $\mathcal{G}_{k,d}$. If $x \in S^{d-1}$, then*

$$P_x = \frac{1}{\alpha} \sum_{j=1}^n \omega_j \|P_{V_j}(x)\|^2 P_{V_j} - \frac{\beta}{\alpha}\,\mathrm{Id}, \tag{11}$$

where $\alpha = \frac{2k(d-k)}{d(d+2)(d-1)}$ and $\beta = \frac{k(kd+k-2)}{d(d+2)(d-1)}$.

Proof. For any $x, y \in S^{d-1}$, the function $V \mapsto \langle P_x, P_V\rangle\langle P_y, P_V\rangle$ belongs to $\mathrm{Pol}_{\leq 4}(\mathcal{G}_{k,d})$. Applying the cubature formula yields

$$\sum_{j=1}^n \omega_j \langle P_x, P_{V_j}\rangle\langle P_y, P_{V_j}\rangle = \int_{\mathcal{G}_{k,d}} \langle P_x, P_V\rangle\langle P_y, P_V\rangle d\sigma_k(V). \tag{12}$$

The function

$$G : (\mathbb{R}x, \mathbb{R}y) \mapsto \int_{\mathcal{G}_{k,d}} \langle P_x, P_V\rangle\langle P_y, P_V\rangle d\sigma_k(V) \tag{13}$$

belongs to $L^2(\mathcal{G}_{1,d} \times \mathcal{G}_{1,d})$ and is zonal. For each variable, it has the form $\mathbb{R}x \mapsto \langle P_x, A(y)\rangle$, where $A(y) = \int_{\mathcal{G}_{k,d}} \langle P_y, P_V\rangle P_V d\sigma_k(V)$, and $\mathbb{R}y \mapsto \langle P_y, A(x)\rangle$, respectively. Since $A(y)$ is self-adjoint and hence a linear combination of projections, $G(\cdot, y)$ and $G(x, \cdot)$ belong to $\mathrm{Pol}_{\leq 2}(\mathcal{G}_{1,d})$. The zonal functions on the projective space are polynomials in the variable $\langle P_x, P_y\rangle = (x, y)^2$, so that G must be of the form $\alpha(x, y)^2 + \beta$. Thus, (12) yields

$$\sum_{j=1}^n \omega_j \langle P_x, P_{V_j}\rangle\langle P_y, P_{V_j}\rangle = \alpha\langle P_x, P_y\rangle + \beta\langle \mathrm{Id}, P_y\rangle. \tag{14}$$

Since (14) holds for every y, we derive

$$\sum_{j=1}^n \omega_j \langle P_x, P_{V_j}\rangle P_{V_j} = \alpha P_x + \beta\,\mathrm{Id}\,. \tag{15}$$

Taking traces in (15) leads to $k\sum_{j=1}^n \omega_j\langle P_x, P_{V_j}\rangle = \alpha + d\beta$, and the property of tight 1-fusion frames gives $\sum_{j=1}^n \omega_j\langle P_x, P_{V_j}\rangle = A_1 = k/d$, so we obtain

$$\alpha + d\beta = k^2/d. \tag{16}$$

Taking $x = y$ in (14) implies $\sum_{j=1}^{n} \omega_j \langle P_x, P_{V_j} \rangle^2 = \alpha + \beta$, and the tight 2-fusion frame property leads to $\sum_{j=1}^{n} \omega_j \langle P_x, P_{V_j} \rangle^2 = A_2 = k(k+2)/(d(d+2))$, so that we obtain

$$\alpha + \beta = k(k+2)/(d(d+2)). \tag{17}$$

Solving for α and β in (16) and (17) yields the required identity. $\square$

To summarize, we have proved:

Theorem 4.3. *Let $\{(V_j, \omega_j)\}_{j=1}^{n}$ be a tight p-fusion frame that is also a cubature of strength 4 for $\mathcal{G}_{k,d}$. For $x \in S^{d-1}$, Algorithm 1 outputs a list L of at most $2p!$ elements of S^{d-1} containing x.*

Algorithm 1 List reconstruction

```
Input: $\{t_j := \|P_{V_j}(x)\|^2\}_{j=p+1}^{n}$.
Output: $L$, $x \in L$.
1: Initialize $L = \emptyset$.
2: Compute the set $\mathcal{S}$ of solutions of the algebraic system of equations in the unknowns $T_1, \dots, T_p$:
       $$\sum_{j=1}^{p} \omega_j T_j^{\ell} = \frac{(k/2)_\ell}{(d/2)_\ell} - \sum_{j=p+1}^{n} \omega_j t_j^{\ell}, \quad 1 \le \ell \le p. \tag{S}$$
3: For every $(t_1, \dots, t_p) \in \mathcal{S}$, and $\alpha, \beta$ defined in Proposition 4.2, compute
       $$P = \frac{1}{\alpha} \sum_{j=1}^{n} \omega_j t_j P_{V_j} - \frac{\beta}{\alpha} \operatorname{Id}.$$
4: If $P$ is a projection of rank 1, compute a unit vector $\xi$ spanning its image and add $\pm\xi$ to $L$.
5: return $L$
```

Note that the actual output list L can be much shorter than $2p!$. In the first place, we can exclude those solutions of (S) that are not real or have negative entries. Moreover, in step 4 one can expect that, for most solutions of (S), the symmetric operator P is not a projection of rank one. Also, the solutions $(t_1, \dots, t_p)$ of $\mathcal{S}$ that do not satisfy $|t_i^{1/2} - t_j^{1/2}|^2 \le \|P_{V_i} - P_{V_j}\|^2$, for every $i < j$, can be removed after step 3 because they violate the consistency conditions

$$\big|\|P_{V_i}(x)\| - \|P_{V_j}(x)\|\big|^2 \le \|P_{V_i}(x) - P_{V_j}(x)\|^2 \le \|P_{V_i} - P_{V_j}\|^2,$$

where $\|P_{V_i} - P_{V_j}\|$ denotes the strong operator norm of $P_{V_i} - P_{V_j}$.

Remark 4.4. 1. For $p = 2$, the assumptions in Theorem 4.3 reduce to $\{(V_j, \omega_j)\}_{j=1}^{n}$ being a cubature of strength 4 for $\mathcal{G}_{k,d}$. Even for $k = 1$, our result extends [6] since we only need $n-2$ elements of the collection $\{\|P_{V_j}(x)\|^2\}_{j=1}^n$ as opposed to all n elements in [6]. This additional flexibility is not for free: We must assume that $x \in S^{d-1}$, and, instead of the two possibilities $\pm x$ in [6], we obtain a list L of 4 elements, one of which is x.

2. Since any symmetric matrix $X \in \mathbb{R}^{d\times d}$ can be written as a sum of weighted orthogonal projectors, (11) can be extended to

$$X = \frac{1}{\alpha}\sum_{j=1}^{n} \omega_j \langle X, P_{V_j}\rangle P_{V_j} - \frac{\beta}{\alpha}\operatorname{trace}(X)\,\mathrm{Id}\,. \tag{18}$$

For $x \in \mathbb{R}^d$ and $X = xx^*$, the tight-1 fusion frame property yields $\operatorname{trace}(X) = \|x\|^2 = \frac{d}{k}\sum_{j=1}^n \omega_j \|P_{V_j}x\|^2$, so that the entire right-hand side of (18) can be computed and hence $\pm x$ can be recovered. Moreover, if V is a random subspace, uniformly distributed in $\mathcal{G}_{k,d}$, i.e., according to σ_k, then the proof of Proposition 4.2 yields that

$$\mathbb{E}(\frac{1}{\alpha}\langle X, P_V\rangle P_V - \frac{\beta}{\alpha}\operatorname{trace}(X)\,\mathrm{Id}) = X, \tag{19}$$

for all symmetric $X \in \mathbb{R}^{d\times d}$. Thus, if $\{V_j\}_{j=1}^n \subset \mathcal{G}_{k,d}$ are independent copies of V, then the law of large numbers implies

$$\frac{1}{n}\sum_{j=1}^{n}\frac{1}{\alpha}\sum_{j=1}^{n}\langle X, P_{V_j}\rangle P_{V_j} - \frac{\beta}{\alpha}\operatorname{trace}(X)\,\mathrm{Id} \to X \quad \text{almost surely.} \tag{20}$$

However, n must be chosen large to obtain an accurate representation of X. In Section 7, we shall see that the random choice of subspaces can be efficient when the algebraic reconstruction formula is replaced with a semidefinite program.

5. Replacing Step 2 with semidefinite programming

We assume in Proposition 4.2 that the weighted subspaces form a cubature of strength 4 for $\mathcal{G}_{k,d}$. However, any real cubature of strength 4 requires at least $\frac{1}{2}d(d+1)$ subspaces, see [22]. Hence, the cardinality scales at least quadratically in the dimension d of the ambient space. In this section, we replace the algebraic reconstruction formula with a semidefinite program similar to the approach in [13], where the case $k=1$ was discussed.

Let $\mathscr{H}$ denote the collection of symmetric matrices in $\mathbb{R}^{d\times d}$. For $\{V_j\}_{j=1}^n \subset \mathcal{G}_{k,d}$, we define the operator

$$\mathcal{F}_n : \mathscr{H} \to \mathbb{R}^n, \quad X \mapsto \frac{d}{k}(\langle X, P_{V_j}\rangle)_{j=1}^n. \tag{21}$$

For $x \in S^{d-1}$, let $f := \frac{d}{k}(\|P_{V_j}(x)\|^2)_{j=1}^n = \mathcal{F}_n(P_x) \in \mathbb{R}^n$, and we now aim to reconstruct x from f. By assuming that the union of the subspaces $\{V_j\}_{j=1}^n$

spans $\mathbb{R}^d$, clearly, P_x is a solution of

$$\min_{X \in \mathscr{H}} (\operatorname{rank}(X)), \quad \text{subject to} \quad \mathcal{F}_n(X) = f, \; X \succeq 0. \tag{22}$$

The notation $X \succeq 0$ stands for X being positive semidefinite. Rank minimization is in general NP-hard. Therefore, (22) is commonly replaced with

$$\min_{X \in \mathscr{H}} (\operatorname{trace}(X)), \quad \text{subject to} \quad \mathcal{F}_n(X) = f, \; X \succeq 0, \tag{23}$$

a semidefinite program, for which efficient solvers such as interior point methods are available on the NEOS Server [21]. For $k = 1$, there is a constant $c > 0$, such that the random choice of at least $cd \log(d)$ subspaces yields that, with high probability, the solutions to (22) and (23) coincide and are unique, cf. [13]. Recently, the logarithmic factor was removed in [12], so that the cardinality of subspaces scales linearly in d. Here, we extend the result to $k > 1$:

Theorem 5.1. *There are constants $c_1, c_2 > 0$ such that, if $x \in \mathbb{R}^d$, $n \geq c_1 d$ and $\{V_j\}_{j=1}^n \subset \mathcal{G}_{k,d}$ are chosen independently identically distributed according to σ_k, then with probability $1 - e^{-c_2 \frac{n}{d}}$ the matrix xx^* is the unique solution to* (23)*.*

To verify Theorem 5.1, we shall first derive deterministic conditions serving equivalence of both problems (21) and (22). Later, we shall verify that these conditions are satisfied with high probability when the subspaces are chosen in the appropriate random fashion. We follow [16] and define, for $X \in \mathscr{H}$,

$$\mathcal{T}(X) = \{Z - X : Z \in \mathscr{H}, \;\; Z \succeq 0 \;\; \operatorname{trace}(Z) \leq \operatorname{trace}(X)\}.$$

A direct computation yields a result stated in [16]:

Proposition 5.2. *The matrix $\hat{X}$ is the unique solution to* (23) *if and only if* $\operatorname{null}(\mathcal{F}_n) \cap \mathcal{T}(\hat{X}) = \{0\}$*.*

While Proposition 5.2 is sufficient and necessary, we also prepare for another sufficient condition that appears easier to check for random subspaces. Let $e = (1, 0, \ldots, 0)^\top \in \mathbb{R}^d$ and T be the linear subspace $\{ey^* + ye^* : y \in \mathbb{R}^d\} \subset \mathscr{H}$. For $Z \in \mathbb{R}^{d \times d}$, denote Z_T its orthogonal projection onto T and $Y_{T^\perp}$ its orthogonal projection onto the orthogonal complement of T. Standard duality arguments from linear programming yield the following observations: suppose that there exists Y be in the range of $\mathcal{F}_n^*$ and consider the conditions

(D-a) $I - Y \succeq 0$ and $Y_T = ee^*$,

(D-b) the right lower $(d-1) \times (d-1)$ block of $I_{T^\top} - Y_{T^\top}$ is positive definite.

If (D-a) is satisfied, then ee^* is a solution to (23), and if in addition (D-b) is satisfied, then this solution is unique. Note that Y is called a *dual certificate* in [13] and that (D-b) and $Y_T = ee^*$ already imply $I - Y \succeq 0$.

Next, we weaken Condition (D-a) and to compensate, we slightly strengthen (D-b) and add a near isometry condition on $\mathcal{F}_n$ just as in [13]. The proof of the following theorem uses Proposition 5.2 and generalizes findings in [13]. Here, $\| \cdot \|_1$ denotes the nuclear norm and $\| \cdot \|_\infty$ the operator norm:

Theorem 5.3. *Let* $\{V_j\}_{j=1}^n \subset \mathcal{G}_{k,d}$ *and* $b = (\|P_{V_j}(e)\|^2)_{j=1}^n$. *Suppose that the following three points are satisfied, where* $0 < r, u_k, \gamma < 1$ *are fixed numbers such that* $\gamma < \frac{u_k}{2}\frac{1-r}{1+r}\frac{1}{\sqrt{1+s^2}}$ *and* $s = \frac{r(u_k+1)+(1-u_k)}{1+r}$.

(1) *For all positive semidefinite matrices* $X \in \mathscr{H}$,

$$\frac{1}{n}\|\mathcal{F}_n(X)\|_{\ell_1} \le (1+r)\|X\|_1. \tag{24}$$

(2) *For all* $X \in T$,

$$\frac{1}{n}\|\mathcal{F}_n(X)\|_{\ell_1} \ge u_k(1-r)\|X\|_\infty. \tag{25}$$

(3) *There exists* Y *in the range of* $\mathcal{F}_n^*$ *such that*

$$\|Y_T - P_e\|_{HS} \le \gamma, \quad \|Y_{T^\perp}\|_\infty \le \frac{1}{2}. \tag{26}$$

Then P_e *is the unique minimizer to* (23).

The Theorem 5.3 was formulated in [13] for $k = 1$, $r = 1/9$, $u_1 = 0.94$, and $\gamma = 1/3$ only. Since $0 < s < 1$, the choice $\gamma = \frac{u_k}{2\sqrt{2}}\frac{1-r}{1+r}$, for instance, is admissible. We can straightforwardly follow the lines of the proof in [13] while keeping track of the constants, so we omit the explicit proof here. It is noteworthy though that it relies on Proposition 5.2 and that the special form of $\mathcal{F}_n$ is not used, so that the statement holds for any linear map from $\mathscr{H}$ to $\mathbb{R}^n$ satisfying the conditions of Theorem 5.3.

Condition (26) is the modification of the above Conditions (D-a) and (D-b), and (24) with (25) are used to make this modification work. Note that (24) is natural in the setting of random subspaces. Assuming that $\{V_j\}_{j=1}^n$ is a tight 1-fusion frame, we obtain, for positive semidefinite $X \in \mathscr{H}$,

$$\|\mathcal{F}_n(X)\|_{\ell_1} = \frac{d}{k}\sum_{j=1}^n \langle X, P_{V_j}\rangle = \sum_{i=1}^d \lambda_i \frac{d}{k}\sum_{j=1}^n \langle P_{x_i}, P_{V_j}\rangle = \sum_{i=1}^d \lambda_i = \|X\|_1,$$

where $\{\lambda_i\}_{i=1}^d$ are the eigenvalues of X and $\{x_i\}_{i=1}^d$ the associated unit norm eigenvectors. We shall verify later that the random choice of subspaces approximates a tight 1-fusion frame, so that (24) is satisfied with high probability. The corresponding inequality from below then holds too but (25) asks for the operator norm in place of the nuclear norm and must hold on T. To compensate, we allow for an additional constant u_k in (25).

In the subsequent sections, we shall verify that the condition of Theorem 5.3 are satisfied with high probability when the subspaces $\{V_j\}_{j=1}^n$ are selected at random.

5.1. Nuclear norm estimates on $\|\mathcal{F}_n(X)\|_{\ell_1}$ for $X \succeq 0$

We shall first verify that $\mathcal{F}_n$ is close to an isometry with high probability:

Theorem 5.4. *For $0 < r < 1$ fixed, there are constants $c(r), C(r) > 0$, such that, for all positive semidefinite matrices X and $n \geq c(r)d$,*

$$(1-r)\|X\|_1 \leq \frac{1}{n}\|\mathcal{F}_n(X)\|_{\ell_1} \leq (1+r)\|X\|_1. \tag{27}$$

holds with probability at least $1 - e^{-C(r)n}$.

To prove Theorem 5.4, we need to extend a result in [48] on the smallest and largest singular values $s_{\min}(P)$ and $s_{\max}(P)$, respectively, in which P has independent sub-exponential rows. Here, we consider independent blocks but there may be dependent rows within each block:

Proposition 5.5. *Let $P := \sqrt{\frac{d}{k}}\big(P_{V_1}, \ldots, P_{V_n}\big)^*$ be a random matrix, in which $\{V_j\}_{j=1}^n$ are identically and independently distributed according to σ_k on $\mathcal{G}_{k,d}$. Then, for every $t \geq 0$, we have with probability at least $1 - 2\exp(-ct^2)$*

$$\sqrt{n} - C\sqrt{d} - t \leq s_{\min}(P) \leq s_{\max}(P) \leq \sqrt{n} + C\sqrt{d} + t,$$

where $c, C > 0$ are absolute constant.

The proof of Proposition 5.5 is given in Appendix A.

Proof of Theorem 5.4. Since any positive semidefinite matrix X can be written by means of its projectors on eigenspaces, it is sufficient to verify

$$1 - r \leq \frac{1}{n}\|\mathcal{F}_n(xx^*)\|_{\ell_1} \leq 1 + r, \quad \forall x \in S^{d-1},$$

in place of (27). We observe that $\|Px\|^2 = \|\mathcal{F}_n(xx^*)\|_{\ell_1}$, so that

$$s_{\min}^2(P) \leq \|\mathcal{F}_n(xx^*)\|_{\ell_1} \leq s_{\max}^2(P)$$

holds. First, we take care of the upper bound. According to Proposition 5.5, we have

$$\frac{1}{n}\|\mathcal{F}_n(P_x)\|_{\ell_1} \leq \frac{1}{n}s_{\max}^2(P) \leq (1 + \frac{1}{\sqrt{n}}(C\sqrt{d} + t))^2,$$

with probability at least $1 - 2e^{-ct^2}$. Choose $\varepsilon > 0$ such that $r/4 = \varepsilon^2 + \varepsilon$ and observe that $\varepsilon \geq \frac{r}{5}$, so that $n \geq c_1 r^{-2} d$ with $c_1 = 25C^2$ implies $n \geq \varepsilon^{-2}C^2 d$. For $t = \sqrt{n}\varepsilon$, we obtain that

$$\frac{1}{\sqrt{n}}s_{\max} \leq (1 + 2\varepsilon)$$

holds with probability at least $1 - 2e^{-cn\varepsilon^2}$. Hence, we have $\frac{1}{n}s_{\max}^2 \leq (1+r)$ with the same probability. Since $\varepsilon^2 \geq r^2/25$, we can adjust c_1 such that $n \geq \frac{25}{cr^2}\ln(2)$ so that $c_2 > 0$ exists and the required upper estimate holds with probability $1 - e^{-c_2 r^2 n}$, The lower estimate can be derived in an analogous way. $\square$

5.2. Operator norm estimates on $\|\mathcal{F}_n(X)\|_{\ell_1}$ for symmetric rank-2 matrices

We shall verify a lower estimate on $\|\mathcal{F}_n(X)\|_{\ell_1}$ involving the operator norm when X is a symmetric rank-2 matrix:

Theorem 5.6. *There is a constant $u_k > 0$, only depending on k, such that the following holds: for $0 < r < 1$ fixed, there exist constants $c(r), C(r) > 0$, such that, for any symmetric rank-2 matrix X, $n \geq c(r)d$,*

$$\frac{1}{n}\|\mathcal{F}_n(X)\|_{\ell_1} \geq u_k(1-r)\|X\|_\infty$$

with probability at least $1 - e^{-C(r)n}$.

We need two preparatory results:

Proposition 5.7. *There is a constant $u_k > 0$ such that, for all $-1 \leq t \leq 1$,*

$$\frac{d}{k}\mathbb{E}\big|\|P_V(e_1)\|^2 - t\|P_V(e_2)\|^2\big| \geq u_k.$$

Lemma 5.8. *The random variable*

$$\xi = \frac{d}{k}|\|P_V(e_1)\|^2 - t\|P_V(e_2)\|^2| \quad \textit{satisfies} \quad \sup_{p\geq 1} 1/p(\mathbb{E}|\xi|^p)^{1/p} \leq 2.$$

Both, Proposition 5.7 and Lemma 5.8, are proven in Appendix B.

Proof of Theorem 5.6. It is sufficient to consider $\|X\|_\infty = 1$, so that $X = P_u - tP_w$, where $u, w \in S^{d-1}$ and $u \perp w$ and $t \in [-1, 1]$. Since σ_k is invariant under $O(\mathbb{R}^d)$, we can assume that u, w are the first two unit vectors e_1 and e_2. We observe

$$\frac{1}{n}\|\mathcal{F}_n(X)\|_1 = \frac{1}{n}\sum_{j=1}^{n}\frac{d}{k}|\|P_{V_j}(e_1)\|^2 - t\|P_{V_j}(e_2)\|^2| =: \frac{1}{n}\sum_{j=1}^{n}\xi_j,$$

where ξ_j are independent copies of

$$\xi = \frac{d}{k}|\|P_V(e_1)\|^2 - t\|P_V(e_2)\|^2|.$$

The Bernstein inequality for sub-exponential random variables as stated in [48] can be used with Lemma 5.8 to obtain

$$\mathbb{P}(|\frac{1}{n}\|\mathcal{F}_n(X)\|_1 - \mathbb{E}\xi| \geq \varepsilon) \leq 2\exp(-cn\min(\frac{\varepsilon^2}{4}, \frac{\varepsilon}{2})),$$

where $c > 0$ is an absolute constant. For $\varepsilon < 2$, we derive with Proposition 5.7

$$\frac{1}{n}\|\mathcal{F}_n(X)\|_1 \geq u_k - \varepsilon,$$

with probability at least $1 - 2\exp(-C_1 n\varepsilon^2)$, where $C_1 = c/4$. The choice $\varepsilon = u_k r$ establishes the required estimate at least for fixed $X \in T$ with probability at least $1 - 2\exp(-C_2 nr^2)$, where $C_2 = C_1 u_k^2$. The remaining part of the proof is the same covering argument as in [13], so we omit this.

$\square$

5.3. Dual certificate: Y_T

It is not hard to see that $\mathcal{F}_n^*$ is given by

$$\mathcal{F}_n^* : \mathbb{R}^n \to \mathscr{H}, \quad (v_j)_{j=1}^n \mapsto \frac{d}{k}\sum_{j=1}^n v_j P_{V_j},$$

so that $\mathcal{F}_n^*\mathcal{F}_n(X) = \frac{d^2}{k^2}\sum_{j=1}^n \langle X, P_{V_j}\rangle P_{V_j}$. If V is a random subspace in $\mathcal{G}_{k,d}$ distributed according to σ_k, then (19) yields, for all $X \in \mathscr{H}$,

$$\frac{d^2}{k^2}\mathbb{E}\langle X, P_V\rangle P_V = aX + b\,\mathrm{trace}(X)I, \tag{28}$$

where $a = \frac{2d(d-k)}{k(d+2)(d-1)}$ and $b = \frac{d(kd+k-2)}{k(d+2)(d-1)}$. As pointed out in (20), the law of large numbers implies that $\mathcal{F}_n^*\mathcal{F}_n$ tends to $\mathcal{S}$, where

$$\mathcal{S} : \mathscr{H} \to \mathscr{H}, \quad X \mapsto aX + b\,\mathrm{trace}(X)I.$$

We can compute its inverse as

$$\mathcal{S}^{-1} : \mathscr{H} \to \mathscr{H}, \quad X \mapsto \frac{1}{a}X - \frac{b}{da}\,\mathrm{trace}(X)I,$$

where the special form of a and b is used. Thus, $\mathcal{F}_n^*\mathcal{F}_n\mathcal{S}^{-1}$ tends to the identity, which motivates the choice of a candidate for Y as

$$\begin{aligned} Y = \frac{1}{n}\mathcal{F}_n^*\mathcal{F}_n\mathcal{S}^{-1}(ee^*) &= \frac{1}{n}\frac{d^2}{k^2}\sum_{j=1}^n \langle \mathcal{S}^{-1}(ee^*), P_{V_j}\rangle P_{V_j} \\ &= \frac{1}{n}\sum_{j=1}^n \big(\frac{d}{ka}\|P_{V_j}(e)\|^2 - \frac{b}{a}\big)\frac{d}{k}P_{V_j}. \end{aligned}$$

In view of tail bound estimates, it will be advantageous though to use an additional cut-off:

Theorem 5.9. *Given $\gamma > 0$, let $\{V_j\}_{j=1}^n$ be identical and independently distributed according to σ_k on $\mathcal{G}_{k,d}$ and define $Y_j = (\frac{d}{ka}\|P_{V_j}(e)\|^2 - \frac{b}{a})\frac{d}{k}P_{V_j}$. Let $Y = \frac{1}{n}\sum_{j=1}^n Y_j 1_{E_j}$, where*

$$E_j = \{\sqrt{\frac{d}{k}}\|P_{V_j}(e)\| \le 2\beta_\gamma\} \tag{29}$$

and $\beta_\gamma > 0$ a sufficiently large constant. Then Y is clearly in the range of $\mathcal{F}_n^*$, and there are constants $c_\gamma, C_\gamma > 0$ such that, for $n \ge c_\gamma d$,

$$\|Y_T - P_e\|_{HS} \le \gamma \tag{30}$$

holds with probability at least $1 - e^{-C_\gamma n/d}$.

Remark 5.10. The structure of Theorem 5.9 and its proof resemble the lines in [13] but we removed the $\log(d)$ from the constraint in E_j. It enables us to remove the logarithmic factor from the lower bound on the number of measurements. Alternatively, Candes and Li removed this factor for $k = 1$ using a slightly different construction in [12].

The proof of Theorem 5.9 is based on two propositions:

Proposition 5.11. *Under the notation of Theorem 5.9, let V be a random subspace distributed according to σ_k on $\mathcal{G}_{k,d}$. Then we have, for all $\beta_\gamma \geq 1/2$,*

$$\mathbb{P}(\sqrt{\frac{d}{k}}\|P_V(e)\| > 2\beta_\gamma) \leq 2e^{k/2}e^{-k\beta_\gamma}. \tag{31}$$

Proposition 5.11 immediately follows from Vershynin's lecture notes on non-asymptotic random matrix theory, (Lemma 9 in Lecture 4 on dimension reduction).

Proposition 5.12. *Under the notation of Theorem 5.9 and $n \geq c_\gamma d$, let*

$$y_j = \xi\frac{d}{k}P_{V_j}e - e, \qquad \xi_j = \frac{d}{ka}\|P_{V_j}(e)\|^2 - \frac{b}{a}. \tag{32}$$

Then we obtain, with probability $1 - e^{-C_\gamma n/d}$,

$$\|\frac{1}{n}\sum_{j=1}^{n} y_j 1_{E_j}\| \leq \frac{\gamma}{2\sqrt{2}}.$$

We shall verify Proposition 5.12 in Appendix C.

Proof of Theorem 5.9. Since $\|Y_T - P_e\|_{HS} \leq \sqrt{2}\|Y_T e - e\|$, we shall use

$$Y_T e - e = \frac{1}{n}\sum_{j=1}^{n} y_j 1_{E_j} - \frac{1}{n}\sum_{j=1}^{n} e 1_{E_j^c}, \tag{33}$$

where y_j is defined by (32) and E_j^c denotes the complement of the set E_j in the appropriate probability space. We shall estimate both terms at the right-hand side of (33) separately. Let us denote

$$\pi(\beta_\gamma) := \mathbb{P}(\sqrt{\frac{d}{k}}\|P_V(e)\| > 2\beta_\gamma).$$

We apply Hoeffding's inequality to obtain

$$\mathbb{P}(|\frac{1}{n}\sum_{j=1}^{n} 1_{E_j^c} - \pi(\beta_\gamma)| \geq t) \leq 2\exp(-2nt^2), \quad \forall\, t > 0. \tag{34}$$

According to Proposition 5.11, we can choose β_γ sufficiently large such that $\pi(\beta_\gamma) \leq \frac{\gamma}{4\sqrt{2}}$. For $t = \frac{1}{4\sqrt{2}}\gamma$, the estimate in (34) implies that

$$\|\frac{1}{n}\sum_{j=1}^{n} e 1_{E_j^c}\| = |\frac{1}{n}\sum_{j=1}^{n} 1_{E_j^c}| \leq \frac{\gamma}{2\sqrt{2}}$$

holds with probability $1 - 2e^{-\frac{\gamma^2}{16}n}$. Combining the latter with Proposition 5.12 yields the estimate (30) with the required probability. $\square$

5.4. Dual certificate: $Y_{T^\top}$

Recall that $Y = \frac{1}{n}\sum_{j=1}^{n} Y_j 1_{E_j}$ with $Y_j = \frac{1}{a}\big(\frac{d}{k}\|P_{V_j}(e)\|^2 - b\big)\frac{d}{k}P_{V_j}$ and $E_j = \{\sqrt{\frac{d}{k}}\|P_{V_j}(e)\| \leq 2\beta_\gamma\}$ were defined in Theorem 5.9. This section is dedicated to verifying the following theorem:

Theorem 5.13. *There are constants $c, C > 0$ such that, for $n \geq cd$,*

$$\|Y_{T^\perp}\|_\infty \leq \frac{1}{2} \tag{35}$$

with probability at least $1 - e^{-Cn}$.

The random variable $\|P_V(x)\|^2$ does not depend on the choice of $x \in S^{d-1}$ and is beta distributed with parameters $(\frac{k}{2}, \frac{d-k}{2})$. Its moments are given by

$$\mathbb{E}\|P_V(x)\|^{2p} = \frac{(k/2)_p}{(d/2)_p} = \frac{k(k+2)\cdots(k+2p-2)}{d(d+2)\cdots(d+2p-2)}, \tag{36}$$

which coincide with the tight p-fusion frame bounds (6) when the weights are constant. Equation (36) is the first step to verify the following lemma whose proof is given in Appendix D:

Lemma 5.14. *If V is a random subspace in $\mathcal{G}_{k,d}$, $k \geq 2$, distributed according to σ_k, then we have, for any $u \in S^{d-1}$ and any integer $p \geq 1$,*

$$\frac{d^p}{k^p}\sqrt{\mathbb{E}\|P_V(u)\|^{4p}} \leq 2^p p!.$$

We need an additional lemma involving nets. A (sperical) ε-net $\mathcal{N}_\varepsilon$ is a finite subset of S^{d-1} such that to any element $x \in S^{d-1}$, there is an element in $\mathcal{N}_\varepsilon$ at distance less than or equals ε.

Lemma 5.15 ([48, Lemma 5.4])**.** *Let $A \in \mathbb{R}^{d\times d}$ be symmetric, and let $\mathcal{N}_\varepsilon$ be an ε-net of S^{d-1} for some $\varepsilon \in [0, \frac{1}{2})$. Then*

$$\|A\|_\infty = \sup_{x\in S^{d-1}} |\langle Ax, x\rangle| \leq (1-2\varepsilon)^{-1} \sup_{x\in\mathcal{N}_\varepsilon} |\langle Ax, x\rangle|.$$

Proof of Theorem 5.13. We see that $Y_{T^\perp} = \frac{1}{n}\sum_{j=1}^n X_j 1_{E_j}$, where E_j is given by (29) and X_j are independent copies of

$$X \sim \frac{1}{a}\big(\frac{d}{k}\|P_V(e)\|^2 - b\big)\frac{d}{k}(P_V)_{T^\perp}.$$

In order to verify (35) with high-probability, we shall show that $\|\frac{1}{n}\sum_{j=1}^n \bar{X}_j\| \leq \frac{1}{2}$ holds with the required probability, where $\bar{X}_j = X_j 1_{E_j}$. We first write

$$\|\sum_{j=1}^n \bar{X}_j\| = \sup_{u\in S^{d-1},\, u\perp e}\big|\sum_{j=1}^n \langle u, \bar{X}_j u\rangle\big| = \sup_{u\in S^{d-1},\, u\perp e}\big|\sum_{j=1}^n \eta_j 1_{E_j}\big|, \tag{37}$$

where η_j are independent copies of

$$\eta = \big(\frac{d}{ka}\|P_V(e)\|^2 - \frac{b}{a}\big)\frac{d}{k}\langle u, (P_V)_{T^\perp}(u)\rangle.$$

Note that we shall take care of the dependency of η on u with a covering argument at the end of the proof. In order to apply a Bernstein inequality, we shall estimate, for an integer $p > 0$,

$$\mathbb{E}|\eta 1_E|^p = \mathbb{E}\Big[\frac{1}{a^p}\big(\frac{d}{k}\|P_V(e)\|^2 - b\big)^p \frac{d^p}{k^p}\langle u, (P_V)_{T^\perp}(u)\rangle^p 1_E\Big].$$

The Cauchy Schwartz inequality yields

$$\mathbb{E}|\eta 1_E|^p \leq \sqrt{\mathbb{E}\big[\frac{1}{a^{2p}}\big(\frac{d}{k}\|P_V(e)\|^2 - b\big)^{2p} 1_E\big]}\sqrt{\mathbb{E}\big[\frac{d^{2p}}{k^{2p}}\langle u, (P_V)_{T^\perp}(u)\rangle^{2p} 1_E\big]}. \tag{38}$$

Next, we estimate the two terms on the right-hand side separately. First, we derive, for sufficiently large β_γ, that $|\frac{d}{k}\|P_V(e)\|^2 - b|1_E \leq 4\beta_\gamma^2$, so that

$$\sqrt{\mathbb{E}\big|\frac{d}{k}\|P_V(e)\|^2 - b\big|^{2p} 1_E} \leq (4\beta_\gamma^2)^p. \tag{39}$$

To estimate the second term on the right-hand side of (38), we observe that $\langle u, (P_V)_{T^\perp}(u)\rangle \leq \|P_V(u)\|^2$. We now combine Lemma 5.14 with (39) to estimate (38) by

$$\mathbb{E}|\eta 1_E|^p \leq (\frac{8}{a}\beta_\gamma^2)^p p! = (\frac{8}{a}\beta_\gamma^2)^2 p! (\frac{8}{a}\beta_\gamma^2)^{p-2}.$$

The Bernstein inequality as stated in [13, Theorem 4.4], for instance, then yields, for all $t > 0$,

$$\mathbb{P}\big(|\sum_{j=1}^n \eta_j 1_{E_j} - n\mathbb{E}\eta_j 1_{E_j}| \geq nt\big) \leq 2\exp(-\frac{n^2t^2}{4(\frac{8}{a}\beta_\gamma^2)^2 + \frac{16}{a}\beta_\gamma^2 nt}).$$

Therefore, there is a constant $C_\beta > 0$ such that, for all $t \geq 1/n$,

$$\mathbb{P}\big(|\frac{1}{n}\sum_{j=1}^n \eta_j 1_{E_j} - \mathbb{E}\eta_j 1_{E_j}| \geq t\big) \leq 2\exp(-C_\beta nt). \tag{40}$$

To estimate $|\mathbb{E}\eta_j 1_{E_j}|$, we first observe that $\frac{d}{k}\mathbb{E}P_V = I$ and evaluating (28) at P_e yields

$$\begin{aligned}\mathbb{E}\eta &= \frac{d^2}{ak^2}\mathbb{E}\langle P_V, P_e\rangle\langle (P_V)_{T^\top}u, u\rangle - \frac{bd}{ak}\mathbb{E}\langle (P_V)_{T^\top}u, u\rangle \\ &= \frac{1}{a}\langle (aP_e + bI)_{T^\top}u, u\rangle - \frac{b}{a}\langle I_{T^\top}u, u\rangle = 0.\end{aligned}$$

Since (36) implies $\mathbb{E}\eta_j^2 \leq c$, for some constant $c > 0$, Cauchy-Schwartz and (31) imply

$$|\mathbb{E}\eta_j 1_{E_j}| = |\mathbb{E}\eta_j 1_{E_j^c}| \leq \sqrt{\mathbb{P}(E_j^c)}\sqrt{\mathbb{E}\eta_j^2} \leq \sqrt{c\pi(\beta_\gamma)} \leq \sqrt{c2e^{k/2}e^{-k\beta_\gamma}}.$$

Therefore, (40) yields

$$\mathbb{P}\big(\frac{1}{n}|\sum_{j=1}^n \eta_j 1_{E_j}| \geq t + \sqrt{c2e^{k/2}e^{-k\beta_\gamma}}\big) \leq 2\exp(-C_\beta n t),$$

for all $t \geq 1/n$. If $c_\gamma > 0$ is a sufficiently large constant and β_γ is sufficiently large too, then $n \geq c_\gamma$ and $t = \frac{1}{c_\gamma}$ yield

$$\mathbb{P}\big(\frac{1}{n}|\sum_{j=1}^n \eta_j 1_{E_j}| \geq 1/4\big) \leq 2\exp(-\tilde{C}_\gamma n), \tag{41}$$

where $\tilde{C_\gamma} > 0$ is some constant only depending on γ.

Next, we apply a covering argument to obtain a bound on $\|\sum_{j=1}^n \bar{X}_j\|$ in (37): let $\mathcal{N}_{1/4}$ be a 1/4-net on S^{d-1}, so that Lemma 5.15 implies

$$\|\frac{1}{n}\sum_{j=1}^n \bar{X}_j\| = \sup_{u\in S^{d-1},\, u\perp e}\big|\langle u, \frac{1}{n}\sum_{j=1}^n \bar{X}_j u\rangle\big| \leq 2\sup_{u\in\mathcal{N}_{1/4}}\big|\langle u, \frac{1}{n}\sum_{j=1}^n \bar{X}_j u\rangle\big|.$$

The net $\mathcal{N}_{1/4}$ can be chosen such that $|\mathcal{N}_{1/4}| \leq 9^d$, cf. [13], so that (41) implies

$$\mathbb{P}\big(\|\frac{1}{n}\sum_{j=1}^n \bar{X}_j\| \geq 1/2\big) \leq \mathbb{P}\big(\sup_{u\in\mathcal{N}_{1/4}}\big|\langle u, \frac{1}{n}\sum_{j=1}^n \bar{X}_j u\rangle\big| \geq 1/4\big) \leq 9^d 2\exp(-\tilde{C}_\gamma n).$$

If $c = c_\gamma$ is sufficiently large and $n \geq cd$, then we can transfer the factor $9^d 2$ into a constant $C > 0$, so that

$$\mathbb{P}\big(\|\frac{1}{n}\sum_{j=1}^n \bar{X}_j\| \geq 1/2\big) \leq \exp(-Cn),$$

which concludes the proof of Theorem 5.13. $\square$

5.5. Proof of Theorem 5.1

We can now assemble all of our findings to verify that the conditions in Theorem 5.3 hold with the required probability:

Proof of Theorem 5.1. We fix $r \in (0,1)$ and choose $\gamma = \frac{u_k}{2\sqrt{2}}\frac{1-r}{1+r}$, where $u_k \in (0,1)$ as in Lemma 5.7. Let c_i and C_i, $i = 1, \ldots, 4$, be suitable positive constants. Theorem 5.4 yields that Condition (24) holds with probability of failure at most $e^{-C_1 n}$, for all $n \geq c_1 d$. Theorem 5.6 implies that Condition (25) holds with probability of failure at most $e^{-C_2 n}$, for all $n \geq c_2 d$. According to Theorem 5.9, the first condition in (26) holds with probability of failure at most $e^{-C_3 n/d}$, for all $n \geq c_3 d$. Theorem 5.13 yields that the second condition in (26) is satisfied with probability of failure at most $e^{-C_4 n}$, for all $n \geq c_4 d$.

Finally, there are constants $c, C > 0$ such that, for all $n \geq cd$, we can estimate $\sum_{i=1}^4 e^{-C_i n/d} \leq e^{-Cn/d}$, so that all conditions in Theorem 5.3 are satisfied with probability at least $1 - e^{-Cn/d}$. This implies Theorem 5.1. $\square$

Remark 5.16. Recently, a more general framework for minimization problems of the form (22) and (23) have been considered in [16]. First, replace $\mathcal{F}$ in (21) with $\mathcal{F}(X) = (\langle X, Z_j \rangle)_{j=1}^n$, where $\{Z_j\}_{j=1}^n \subset \mathbb{R}^{d\times d}$ is a collection of independent random matrices with independent identically distributed Gaussian entries. If $x \in S^{d-1}$ and $b = \mathcal{F}(P_x)$, then results in [16] imply that, with high probability, the solution to (23) is unique and also solves (22). These results do not apply in our setting though, because we must deal with orthogonal projectors.

6. Stability under distortions

We shall verify that our approach is robust against contamination of the measurements. For $x \in \mathbb{R}^d$ and following [13], we assume we measure

$$f_j = \|P_{V_j}(x)\|^2 + \eta_j,$$

where $\eta = \{\eta_j\}_{j=1}^n$ is a distortion bounded by $\|\eta\|_{\ell_2} \leq \epsilon$, for some $\epsilon > 0$. We replace (23) with

$$\min_{X \in \mathscr{H}} (\operatorname{trace}(X)) \quad \text{subject to} \quad \|\mathcal{F}_n(X) - f\| \leq \epsilon, \quad X \succeq 0. \tag{42}$$

For $\hat{X}$ being a minimizer of (42), we extract $\pm\hat{x}$ from $\hat{x}\hat{x}^* := \hat{\lambda}_1 P_{v_1}$, where v_1 is the unit norm eigenvector of $\hat{X}$ associated to the largest eigenvalue $\hat{\lambda}_1$.

Theorem 6.1. *There are constants $c_1, c_2, c_3 > 0$ such that, if $x \in \mathbb{R}^d$, $n \geq c_1 d$ and $\{V_j\}_{j=1}^n$ are chosen independently identically distributed according to σ_k on $\mathcal{G}_{k,d}$, then with probability $1 - e^{-c_2 \frac{n}{d}}$ the solution to (42) satisfies*

$$\|x - \varphi\hat{x}\| \leq c_3 \min(\|x\|, \epsilon/\|x\|), \quad \textit{for some } \varphi \in \{\pm 1\}.$$

The proof is derived by following the lines in [13] and keeping track of the constants. For the sake of completeness, we give a brief outline:

Proof. Without loss of generality, we may assume $x = e$ and define $R := ee^* - \hat{X}$. Rescaling and a rotation then yield the general result. Given $0 < r, u_k < 1$, we choose $\gamma < \frac{u_k}{2\sqrt{2}} \frac{1-r}{1+r}$, so that the conditions in Theorem 5.3 are satisfied with probability $1 - e^{-c_2 \frac{n}{d}}$. By following the lines in [13], we derive

$$\begin{aligned} \|R_{T^\perp}\|_1 &\leq 4\epsilon(1+r) + 2\sqrt{2}\gamma \|R_T\|_\infty, \\ u_k(1-r)\|R_T\|_\infty &\leq 2n^{-1/2}\epsilon + (1+r)\|R_{T^\perp}\|_1. \end{aligned} \tag{43}$$

Merging the two inequalities yield

$$\|R_{T^\perp}\|_1 \leq \frac{4u_k(1-r)(1 + r + \frac{\sqrt{2}n^{-1/2}}{u_k(1-r)}\gamma)}{u_k(1-r) - 2\sqrt{2}(1+r)\gamma}\epsilon,$$

and our condition on γ implies that the denominator is positive. Thus, we can define a constant $\tilde{c}_1 > 0$ such that $\|R_{T^\perp}\|_1 \leq \tilde{c}_1\epsilon$. Using (43) yields $\|R_T\|_\infty \leq \tilde{c}_2\epsilon$, for some $\tilde{c}_2 > 0$, so that $\|ee^* - \hat{X}\|_\infty = \|R\|_\infty \leq (\tilde{c}_1 + \tilde{c}_2)\epsilon$. It was verified in [13] that the latter is sufficient to deduce the existence of a constant $c_3 > 0$ satisfying $\|e - \varphi\hat{x}\| \leq c_3 \min(1, \epsilon)$. □

7. Numerical experiments

We shall present some numerical experiments illustrating Theorem 5.1. Let $x \in S^{d-1}$ and observe that $V \in \mathcal{G}_{k,d}$ is uniformly distributed if and only if $P_V = Z(Z^*Z)^{-1}Z^*$ for some $Z \in \mathbb{R}^{d\times k}$ with independent $\mathcal{N}(0,1)$ entries, cf. [17, Theorem 2.2.2]. Thus, we can easily generate pseudo-random orthogonal projectors $\{P_{V_j}\}_{j=1}^n$. Since $\|P_{V_j^\perp}(x)\|^2 + \|P_{V_j}(x)\|^2 = \|x\|^2$, we shall restrict us to $k \leq d/2$. We follow the numerical experiments in [13], where the measurement vector f is given by

$$f_j = \|P_{V_j}(x)\|^2 + \eta_j, \tag{44}$$

and $\eta = \{\eta_j\}_{j=1}^n$ are independent Gaussian random variables with zero mean and variance Σ_η. It then is standard to replace (23) with the relaxation

$$\min_{X\in\mathscr{H}} (\|\mathcal{F}(X) - f\|^2 + \lambda \operatorname{trace}(X)), \quad \text{subject to} \quad X \succeq 0. \tag{45}$$

As in [13], we use the software package Templates for First-Order Conic Solvers (TFOCS) [8] to solve (45). If $\hat{X}$ is the solution to (45), then we define $\pm\hat{x} \in S^{d-1}$ as the normalized eigenvector corresponding to the largest eigenvalue of $\hat{X}$, because we already know that x has unit norm.

First, we follow a numerical test from [11] for the noise free setting. As in [11] for $k = 1$, the computed approximation is visually indistinguishable from the test signal when $k = 10$ and $k = 20$, cf. Fig. 1. We compare the approximation error $\min(\|x \pm \hat{x}\|)$ for several choices of k in a separate experiment in Fig. 2. There, we observe that the error decreases with growing k when λ is kept fixed. An explanation may be that, as k grows, the terms $\{\|P_{V_1}(x)\|^2\}_{j=1}^n$ increase



(a) $k = 10$

(b) $k = 20$

Figure 1: We choose the original signal x uniformly distributed on the sphere S^{d-1} and the measurements b are noise-free. As in [11], where $k = 1$ was used, the approximation is computed for $d = 128$, $n = 6d$, and $\lambda = 0.05$. Here, also for $k > 1$, we see that original and computed signal are visually indistinguishable.



(a) We chose $d = 20$ and $\lambda = 10^{-6}$. The error decreases with growing subspace dimension k and cardinality n. For $k = 1$, the error is above the range of this plot.

(b) For $d = 100$, $n = 10d$, and $\lambda = 10^{-2}$, the error decreases with growing k.

Figure 2: The signal is $x = (1, 0, \dots, 0)^\top \in S^{d-1}$. We computed the error $\min(\|x \pm \hat{x}\|)$ averaged over 100 iterations. Since there is no noise, λ should be small.

so that the ratio between the two competing terms in (45) changes in favor to larger k when λ is held fixed.

Second, we aim to reconstruct the signal x in a noisy setting for several choices of Σ_η. The relaxation parameter λ must still be adopted to the strength of the noise. If Σ_η is indeed known, then λ can be chosen according to the discrepancy principle, but, in practice Σ_η can only be estimated and cross-validation is more commonly used, see [39] for advantages and problems with various parameter estimation techniques. Here, we shall adopt the L-curve criterion [39], see also [29, 44], so that we choose λ according to the maximal curvature of the two-dimensional curve $\big(\|\mathcal{F}(\hat{X}_\lambda) - f\|^2, \operatorname{trace}(\hat{X}_\lambda)\big)$, where $\hat{X}_\lambda$ is the λ-dependent solution to (45).

In Fig. 3, we adjust the noise level Σ_η so that $\mathrm{SNR} = \frac{\Sigma_f}{\Sigma_\eta}$ is fixed when k changes. Here, Σ_f is the standard deviation of the collection $\{\|P_{V_j}(x)\|^2\}_{j=1}^n$.



(a) SNR = 1 (b) SNR = 2



(c) SNR = 4 (d) SNR = 8

Figure 3: For the given data vector b, we fix the SNR. We chose $n = 10d$, and $k = 1, 8, 32$. The original signal x is again a normalized sine wave with $d = 128$ samples.

The three choices of k then seem to perform equally well. Thus, we do not seem to lose anything by using orthogonal projectors whose rank is not one.

It appears reasonable to believe that properties of the signal x that are known a-priori could be used to add further penalty terms in (45) improving the reconstruction.

8. Conclusions

We derived algebraic conditions on subspaces $\{V_j\}_{j=1}^n$ in Proposition 4.2 to reconstruct any signal up to its sign from the norms of its subspace components by means of a closed formula. Those conditions require the existence of nonnegative weights $\{\omega_j\}_{j=1}^n$ such that $\{(V_j, \omega_j)\}_{j=1}^n$ is a cubature for $\mathcal{G}_{k,d}$ of strength 4. The latter extends the results in [6] in two ways: We deal not only with one- but with k-dimensional subspaces, and we do not require the weights to be constant. Next, we tackled the erasure of subspace components using similar algebraic conditions. If $\{(V_j, \omega_j)\}_{j=1}^n$ is in addition a tight p-fusion frame and $x \in S^{d-1}$, then $\{\|P_{V_j}(x)\|^2\}_{j=1}^n$ can essentially be determined from any subset of cardinality $n - p$, see Proposition 4.1. Indeed, the erased p norms can be algorithmically recovered up to permutation. Note that any cubature of strength 4 is also a tight 2-fusion frame so that we only need $n - 2$ norms of subspace components to recover x. To our knowledge, the presented algorithm is even novel for 1-dimensional subspaces.

The cardinality of cubatures of strength 4 scales at least quadratically in the ambient dimension d. For 1-dimensional random subspaces, semidefinite programming was used in [13] to reduce the number of measurements to a linear scale up to a logarithmic factor, and this factor was later removed in [12]. Recently, an optimizationless approach for 1-dimensional subspaces was proposed in [23], but the lower bound on the number of measurements involves the additional logarithmic factor as in [13].

Here, we were dealing with general k-dimensional random subspaces in Section 7 and verified that the signal can be recovered in a stable fashion by using semidefinite programming. It is noteworthy that we avoid the logarithmic factor and reduce the number of required measurements to a linear scale in the ambient dimension.

Acknowledgements

The authors would like to thank Thomas Bauer for the proof of Proposition 4.1 and Pierre Thibault for discussions on diffraction imaging. The authors are also thankful to Frank Valentin and Christian Reiher for valuable discussions on the proof of Theorem 5.1. M. E. is supported by the NIH/DFG Research Career Transition Awards Program (EH 405/1-1/575910).

Appendix A. Proofs for Section 5.1

Our proof of Proposition 5.5 requires a lemma for preparation:

Lemma Appendix A.1 ([48, Lemma 5.36]). *If* $B \in \mathbb{R}^{n\times d}$ *satisfies* $\|B^*B - I\|_\infty \leq \max(\delta, \delta^2)$, *for some* $\delta > 0$, *then*

$$1 - \delta \leq s_{\min}(B) \leq s_{\max}(B) \leq 1 + \delta. \tag{A.1}$$

Conversely, if B satisfies (A.1), *then* $\|B^*B - I\|_\infty \leq 3\max(\delta, \delta^2)$.

Proof of Proposition 5.5. To verify Proposition 5.5, we want to apply Lemma Appendix A.1 with $B = \frac{1}{\sqrt{n}}P$. First, we take care of the upper estimate on $s_{\max}(P)$. We must check that

$$\|\frac{1}{n}\frac{d}{k}\sum_{j=1}^{n} P_{V_j} - I\| \leq \max(\delta, \delta^2) =: \varepsilon, \quad \text{where} \quad \delta = C\sqrt{\frac{d}{n}} + \frac{t}{\sqrt{n}}. \tag{A.2}$$

Let $\mathcal{N}_{1/4}$ be a $\frac{1}{4}$-net, so that we can estimate with Lemma 5.15,

$$\begin{aligned}\|\frac{1}{n}P^*P - I\| &= \|\frac{1}{n}\frac{d}{k}\sum_{j=1}^{n} P_{V_j} - I\| \\ &\leq 2\max_{x\in\mathcal{N}_{1/4}} |\langle(\frac{1}{n}\frac{d}{k}\sum_{j=1}^{n} P_{V_j} - I)x, x\rangle| \\ &= 2\max_{x\in\mathcal{N}_{1/4}} |\frac{1}{n}\|Px\|^2 - 1|.\end{aligned}$$

Thus, we must verify with the required probability that

$$\max_{x\in\mathcal{N}_{1/4}} |\frac{1}{n}\|Px\|^2 - 1| \leq \frac{\varepsilon}{2}. \tag{A.3}$$

To derive this estimate, we define random variables $Z_j = \sqrt{\frac{d}{k}}\|P_{V_j}(x)\|$ so that $\sum_{j=1}^{n} Z_j^2 = \|P(x)\|^2$. Since $\mathbb{E}(Z_j^2) = 1$, we can estimate

$$\begin{aligned}\|Z_j^2\|_{\psi_1} &:= \sup_{p\geq 1} p^{-1}(\mathbb{E}Z_j^{2p})^{1/p} \\ &= \frac{d}{k}\sup_{p\geq 1} p^{-1}(\mathbb{E}\|P_{V_j}(x)\|^{2p})^{1/p} \\ &= \frac{d}{k}\sup_{p\geq 1} p^{-1}(\frac{(k/2)_p}{(d/2)_p})^{1/p} \leq 1,\end{aligned}$$

where the last inequality can be derived by induction over p. According to [48, Remark 5.18], $\|Z_j^2 - 1\|_{\psi_1} \leq 2$, and we obtain from the Bernstein type inequality [48, Corollary 5.17]

$$\begin{aligned}\mathbb{P}(|\frac{1}{n}\sum_{j=1}^{n}\frac{d}{k}\|P_{V_j}(x)\|^2 - 1| \geq \varepsilon/2) &= \mathbb{P}(|\frac{1}{n}\sum_{j=1}^{n} Z_j^2 - 1| \geq \varepsilon/2) \\ &\leq 2\exp(-cn\min(\frac{\varepsilon^2}{16}, \frac{\varepsilon}{4})) \\ &= 2\exp(-n\frac{c}{16}\delta^2) \\ &\leq 2\exp(-\frac{c}{16}(C^2d + t^2)),\end{aligned}$$

where the last line follows from (A.2). Since the net can be chosen such that $|\mathcal{N}_{1/4}| \leq 9^d$, cf. [48], we can verify (A.3) by

$$\begin{aligned}\mathbb{P}\big(\max_{x\in\mathcal{N}_{1/4}} |\frac{1}{n}\sum_{j=1}^{n}\frac{d}{k}\|P_{V_j}(x)\|^2 - 1| \geq \varepsilon/2\big) &\leq 9^d 2\exp(-\frac{c}{16}(C^2d + t^2)) \\ &\leq 2\exp(-\frac{c}{16}t^2),\end{aligned}$$

where we assume $C \geq 4\sqrt{\ln(9)/c}$. The latter does not cause any trouble because c is a constant independent of ε. This finally yields

$$\mathbb{P}(s_{\max}(P) \geq \sqrt{n} + C\sqrt{d} + t) \leq 2\exp(-\frac{c}{16}t^2).$$

The estimates on $s_{\min}(P)$ are derived analoguously. □

Appendix B. Proofs for Section 5.2

We shall prove Proposition 5.7 and Lemma 5.8.

Proof of Proposition 5.7. Since the integral is always nonzero, we only need to take care of the limit $d \to \infty$. We first see that

$$\begin{aligned}\frac{d}{k}\mathbb{E}\big|\|P_V(e_1)\|^2 - t\|P_V(e_2)\|^2\big| &= \frac{d}{k}\int_{\mathcal{G}_{k,d}} \big|\|P_V(e_1)\|^2 - t\|P_V(e_2)\|^2\big| d\sigma_k(V)\\ &= \frac{d}{k}\int_{\mathcal{V}_{2,d}} \big|\sum_{i=1}^{k} m_{i,1}^2 - t\sum_{i=1}^{k} m_{i,2}^2\big| d\nu_2(M),\end{aligned}$$

where $\mathcal{V}_{2,d} = \{M = (m_{i,j}) \in \mathbb{R}^{d\times 2} : M^*M = I\}$ denotes the Stiefel-manifold endowed with the standard probability measure ν_2. If M is a random matrix, distributed according to ν_2, then, according to [25, Proposition 7.5], the upper $k \times 2$ block of M multiplied by d converges in distribution (for $d \to \infty$) towards a random $k \times 2$ matrix whose entries are standard normal i.i.d.. Let us denote the underlying probability measure on $\mathbb{R}^{k\times 2}$ by $\mathcal{N}(0, I_k \otimes I_2)$. The convergence in distribution implies that, for $d \to \infty$,

$$d\mathbb{E}\big|\|P_V(e_1)\|^2 - t\|P_V(e_2)\|^2\big| \to \int_{\mathbb{R}^{k\times 2}} \big|\|N(e_1)\|^2 - t\|N(e_2)\|^2\big| d\mathcal{N}(0, I_k \otimes I_2)(N).$$

Since the right-hand side is bigger than 0, for all $-1 \leq t \leq 1$, compactness and continuity arguments suffice to conclude the statement of the lemma. □

Proof of Lemma 5.8. By using the expressions of $\mathbb{E}\|P_V(x)\|^{2p}$ in (36), we can estimate

$$\begin{aligned}(\mathbb{E}|\xi|^p)^{1/p} &\leq \frac{d}{k}\Big(\mathbb{E}(\|P_V(u)\|^2 + |t|\|P_V(w)\|^2)^p\Big)^{1/p}\\ &= \frac{d}{k}\Big(\sum_{m=0}^{p}\binom{p}{m}\mathbb{E}(\|P_V(u)\|^{2m}|t|^{p-m}\|P_V(w)\|^{2(p-m)})\Big)^{1/p}\\ &\leq \frac{d}{k}\Big(\sum_{m=0}^{p}\binom{p}{m}\sqrt{\mathbb{E}(\|P_V(u)\|^{4m})}\sqrt{\mathbb{E}(\|P_V(w)\|^{4(p-m)})}\Big)^{1/p}\\ &\leq \Big(\sum_{m=0}^{p}\binom{p}{m}\sqrt{\frac{d^2k(k+2)}{k^2d(d+2)}\cdots\frac{d^2(k+4m-4)(k+4m-2)}{k^2(d+4m-4)(d+4m-2)}}\\ &\qquad\cdot\sqrt{\frac{d^2k(k+2)}{k^2d(d+2)}\cdots\frac{d^2(k+4(p-m)-4)(k+4(p-m)-2)}{k^2(d+4(p-m)-4)(d+4(p-m)-2)}}\Big)^{1/p}.\end{aligned}$$

Now, we check that, for $k \geq 2$, $\ell = 0, 1, 2, \ldots$,

$$\frac{d^2}{k^2}\frac{(k+4\ell)}{(d+4\ell)}\frac{(k+4\ell+2)}{(d+4\ell+2)} \leq (1+\frac{4\ell}{k})(1+\frac{4\ell+2}{k}) \leq 2^2(\ell+1)^2. \tag{B.1}$$

Thus, we obtain

$$\begin{aligned}
(\mathbb{E}|\xi|^p)^{1/p} &\leq \Big(\sum_{m=0}^{p}\binom{p}{m}\sqrt{2^{2m}(m!)^2}\sqrt{2^{2(p-m)}((p-m)!)^2}\Big)^{1/p} \\
&\leq \Big(\sum_{m=0}^{p}\binom{p}{m}2^p m!(p-m)!\Big)^{1/p} \\
&\leq 2\Big(p!(p+1)\Big)^{1/p} \leq 2p,
\end{aligned}$$

where the last inequality holds provided that $p \geq 3$. The latter follows from $p!(p+1) \leq p^p$, for $p \geq 3$, which can be shown by induction. Similar arguments as above yield $1/2(\mathbb{E}|\xi|^2)^{1/2} \leq \sqrt{3} \leq 2$ and $\mathbb{E}|\xi|^2) \leq 2$ so that we obtain the required statement. $\square$

Appendix C. Proofs for Section 5.3

Before we can verify Proposition 5.12 using a Bernstein inequality, we require some bounds on the random vectors y_j given in (32) and $\bar{y}_j := y_j 1_{E_j}$:

Lemma Appendix C.1. *There is a constant $c > 0$ such that the following points hold:*

(a) $\mathbb{E}(\|y_j\|^2) \leq cd$,

(b) $\|\bar{y}_j\|_2 \leq c\sqrt{\frac{d}{k}}\beta_\gamma^3$,

(c) $\|\mathbb{E}\bar{y}\|_2 \leq c\sqrt{2e^{k/2}e^{-k\beta_\gamma}}$.

Proof. (a) By applying (36) for $p = 1$ and $p = 2$, a short calculations yields $\mathbb{E}\xi_j\|P_V(e)\|^2 = \frac{k}{d-k}$, and we can compute

$$\begin{aligned}
\mathbb{E}(\|y_j\|^2) &= \mathbb{E}(\|\xi_j\frac{d}{k}P_V e - e\|^2) \\
&= \mathbb{E}(\xi_j^2\frac{d^2}{k^2}\|P_V(e)\|^2 - 2\xi_j\frac{d}{k}\|P_V e\|^2 + 1) \\
&= \frac{d^2}{k^2}\mathbb{E}(\xi_j^2\|P_V(e)\|^2) - 2\frac{d}{d-k} + 1 \\
&= \frac{d^2}{k^2}\mathbb{E}((\frac{d}{ak}\|P_V(e)\|^2 - \frac{b}{a})^2\|P_V(e)\|^2) - 2\frac{d}{d-k} + 1 \\
&= \frac{d^2}{a^2k^2}\mathbb{E}\big((\frac{d^2}{k^2}\|P_V(e)\|^4 - 2\frac{bd}{k}\|P_V(e)\|^2 + b^2)\|P_V(e)\|^2\big) + 1 \\
&= \frac{d^2}{a^2k^2}\Big(\frac{d^2k(k+2)(k+4)}{k^2d(d+2)(d+4)} - 2\frac{bdk(k+2)}{kd(d+2)} + \frac{b^2k}{d}\Big) - 2\frac{d}{d-k} + 1.
\end{aligned}$$

We can estimate a and b by a constant independently of d, which implies (a).

(b) If ξ_j is given by (32), then a direct calculation yields that there is a constant $c > 0$ such that $|\xi_j| \leq c\beta_\gamma^2$ holds on the event E_j. Since $\|y_j\|^2 = \xi_j^2 \frac{d^2}{k^2}\|P_{V_j}(e)\|^2 - 2\xi_j \frac{d}{k}\|P_{V_j}(e)\|^2 + 1$, we can use $\frac{d}{k}\|P_{V_j}(e)\|^2 \leq 4\beta_\gamma^2$ on E_j to obtain the rough estimate

$$\|\bar{y}_j\|^2 \leq c_1 \frac{d}{k}(4\beta_\gamma^2)^3,$$

where $c_1 > 0$ is some constant.

(c) The term $\mathbb{E}P_{V_j}$ is a multiple of the identity, so that the entries of $P_{V_j}(e)$ have zero mean except for the first one. Moreover, (28) yields

$$\frac{d^2}{k^2}\mathbb{E}\|P_{V_j}(e)\|^2 P_{V_j}(e) = (a+b)e$$

and $a + b = \frac{d(k+2)}{(d+2)k}$, which implies that the entries of $y_j = \frac{d^2}{ak^2}\|P_{V_j}(e)\|^2 P_{V_j}(e) - \frac{bd}{ak}P_{V_j}(e) - e$ have zero mean. In other words, $y_j = \frac{d^2}{ak^2}p_{1,1}p_{\ell,1} - \frac{bd}{ak}p_{\ell,1}$, for $\ell = 2, \ldots, d$ have zero mean, where we used $P_{V_j} = (p_{\ell,m})$. The entries $p_{1,1}$ and $p_{\ell,1}$ are even uncorrelated, for $\ell > 1$, so that

$$\mathbb{E}(p_{2,1}) = 0, \quad \mathbb{E}(p_{1,1}p_{2,1}) = 0.$$

We have $-1 \leq p_{2,1} \leq 1$ and its distribution is symmetric around 0 (multiply the first row of P_{V_j} with -1 and the first column too. This is done by $UP_{V_j}U^*$, where U is the identity but the first entry is -1). This way, we see that any value of $p_{1,1}$ corresponds to an entry $p_{2,1}$ and also its negative. Thus, we have $\mathbb{E}(p_{2,1}1_{E_j}) = 0$ and $\mathbb{E}(p_{1,1}p_{2,1}1_{E_j}) = 0$, so that the entries of $\bar{y}_j = y1_{E_j}$ have zero mean except for the first entry. This implies with the identities $\|\mathbb{E}\bar{y}_j\| = |\mathbb{E}\langle y_j - \bar{y}_j, e\rangle| = |\mathbb{E}\langle y_j, e\rangle 1_{E_j^c}|$ and (31) that

$$\|\mathbb{E}\bar{y}_j\| \leq \sqrt{\mathbb{P}(E_j^c)}\sqrt{\mathbb{E}|\langle y_j, e\rangle|^2} \leq \sqrt{2e^{k/2}e^{-k\beta_\gamma}}\sqrt{\mathbb{E}|\langle y_j, e\rangle|^2}. \tag{C.1}$$

Next, we must check that $\mathbb{E}|\langle y_j, e\rangle|^2$ is bounded. If we apply (36), then we obtain that

$$\mathbb{E}|\langle y_j, e\rangle|^2 = \mathbb{E}\big(\frac{d^2}{ak^2}\|P_{V_j}(e)\|^4 - \frac{bd}{ak}\|P_{V_j}(e)\|^2 - 1\big)^2$$

is bounded by a constant, so that (C.1) implies (c). ☐

Proof of Proposition 5.12. The vector Bernstein inequality as stated in [38, Theorem 12] yields, for $\sum_{j=1}^n \mathbb{E}\|y_j\|^2 \leq A$ and for all $t \leq A/\max_{j=1,\ldots,n}\|\bar{y}_j\|$ that

$$\mathbb{P}\big(\|\sum_{j=1}^n \bar{y}_j - n\mathbb{E}\bar{y}\| \geq \sqrt{A} + t\big) \leq \exp(-t^2/4A).$$

We apply our estimates in Lemma Appendix C.1, so that replacing t/n with t yields the following: for $A = cnd$ and for all $t \leq \sqrt{\frac{d}{k}\frac{1}{\beta_\gamma^3}}$,

$$\mathbb{P}\big(\|\frac{1}{n}\sum_{j=1}^{n}\bar{y}_j - \mathbb{E}\bar{y}_j\| \geq \sqrt{c\frac{d}{n}} + t\big) \leq \exp(-nt^2/4cd).$$

Therefore, we also derive

$$\mathbb{P}\big(\|\frac{1}{n}\sum_{j=1}^{n}\bar{y}_j\| \geq \sqrt{c\frac{d}{n}} + t + c\sqrt{2e^{k/2}e^{-k\beta_\gamma}}\big) \leq \exp(-nt^2/4cd).$$

We choose β_γ sufficiently large, t sufficiently small, and c_γ sufficiently large, such that,

$$\sqrt{c\frac{d}{n}} + t + c\sqrt{2e^{k/2}e^{-k\beta_\gamma}} \leq \frac{\gamma}{2\sqrt{2}}, \quad \text{for all } n \geq c_\gamma d,$$

which concludes the proof of Proposition 5.12. □

Appendix D. Proofs for Section 5.4

Proof of Lemma 5.14. By grouping $\frac{(k/2)_{2p}}{(d/2)_{2p}}$ into pairs of two, the estimates (B.1) imply

$$\begin{aligned}\mathbb{E}\frac{d^{2p}}{k^{2p}}\|P_V(u)\|^{4p} &= \frac{d^{2p}}{k^{2p}}\frac{(k/2)_{2p}}{(d/2)_{2p}}\\ &= \frac{d^2k(k+2)}{k^2d(d+2)}\cdot\frac{d^2(k+4)(k+6)}{k^2(d+4)(d+6)}\cdots\frac{d^2(k+4p-4)(k+4p-2)}{k^2(d+4p-4)(d+4p-2)}\\ &\leq 2^{2p}(p!)^2.\end{aligned}$$

□

References

[1] C. Bachoc, *Designs, groups and lattices*, J. Theor. Nombres Bordeaux (2005), 25–44.

[2] ______, *Linear programming bounds for codes in Grassmannian spaces*, IEEE Trans. Inf. Th. **52** (2006), no. 5, 2111–2125.

[3] C. Bachoc, E. Bannai, and R. Coulangeon, *Codes and designs in Grassmannian spaces*, Discrete Mathematics **277** (2004), 15–28.

[4] C. Bachoc, R. Coulangeon, and G. Nebe, *Designs in Grassmannian spaces and lattices*, J. Algebraic Combinatorics **16** (2002), 5–19.

[5] C. Bachoc and M. Ehler, *Tight p-fusion frames*, Appl. Comput. Harmon. Anal., in press, DOI:10.1016/j.acha.2012.07.001 (2012).

[6] R. Balan, B. G. Bodmann, P. G. Casazza, and D. Edidin, *Painless reconstruction from magnitudes of frame coefficients*, J. Fourier Anal. Appl. **15** (2009), no. 4, 488–501.

[7] R. Balan, P. Casazza, and D. Edidin, *On signal reconstruction without phase*, Appl. Comput. Harmon. Anal **20** (2006), 345–356.

[8] S. Becker, E. J. Candès, and M. Grant, *Templates for convex cone problems with applications to sparse signal recovery*, Mathematical Programming Computation **3** (2011), no. 3, 165–218.

[9] J. Bochnak, M. Coste, and M. Roy, *Real algebraic geometry*, Springer-Verlag, 1998.

[10] B. G. Bodmann and V. I. Paulsen, *Frames, graphs and erasures*, Lin. Alg. Appl. **404** (2005), 118–146.

[11] E. J. Candès, Y. Eldar, T. Strohmer, and V. Voroninski, *Phase retrieval via matrix completion*, arXiv:1109.0573v2 (2011).

[12] E. J. Candès and X. Li., *Solving quadratic equations via phaselift when there are about as many equations as unknowns*, ArXiv (2012).

[13] E. J. Candès, T. Strohmer, and V. Voroninski, *PhaseLift: Exact and stable signal recovery from magnitude measurements via convex programming*, arXiv:1109.4499v1, to appear in Communications on Pure and Applied Mathematics.

[14] P. G. Casazza and J. Kovacevic, *Equal-norm tight frames with erasures*, Adv. Comput. Math. **18** (2003), no. 2-4, 387–430.

[15] P. G. Casazza, G. Kutyniok, and S. Li, *Fusion frames and distributed processing*, Appl. Comput. Harmon. Anal. **25** (2008), no. 1, 114–132.

[16] V. Chandrasekaran, B. Recht, P. A. Parrilo, and A. S. Willsky, *The convex geometry of linear inverse problems*, arXiv:1012.0621v3 (2012).

[17] Y. Chikuse, *Statistics on special manifolds*, Lecture Notes in Statistics, Springer, New York, 2003.

[18] O. Christensen, *An Introduction to Frames and Riesz Bases*, Birkhäuser, Boston, 2003.

[19] A. M. Cohen, H. Cuypers, and H. Sterk (eds.), *Some tapas of computer algebra*, Algorithms and Computation in Mathematics, vol. 4, Springer, 1999.

[20] D. Cox, J. Little, and D. O'Shea, *Ideals, Varieties, and Algorithms*, Springer, 1996.

[21] J. Czyzyk, M. Mesnier, and J. Moré, *The NEOS server*, IEEE Computational Science & Engineering **5** (1998), no. 3, 68–75.

[22] P. de la Harpe and C. Pache, *Cubature formulas, geometrical designs, reproducing kernels, and Markov operators*, Infinite groups: geometric, combinatorial and dynamical aspects (Basel), vol. 248, Birkhäuser, 2005, pp. 219–267.

[23] L. Demanet and P. Hand, *Stable optimizationless recovery from phaseless linear measurements*, ArXiv (2012).

[24] J. Drenth, *Principles of protein x-ray crystallography*, Springer, 2010.

[25] M. L. Eaton, *Group invariance applications in statistics*, Regional Conference Series in Probability and Statistics, Institute of Mathematical Statistics, 1989.

[26] M. Ehler, *On multivariate compactly supported bi-frames*, J. Fourier Anal. Appl. **13** (2007), no. 5, 511–532.

[27] M. Ehler and J. Galanis, *Frame theory in directional statistics*, Stat. Probabil. Lett. **81** (2011), no. 8, 1046–1051.

[28] M. Ehler and B. Han, *Wavelet bi-frames with few generators from multivariate refinable functions*, Appl. Comput. Harmon. Anal. **25** (2008), no. 3, 407–414.

[29] M. Ehler and K. Koch, *The construction of multiwavelet bi-frames and applications to variational image denoising*, Int. J. Wavelets, Multiresolut. Inf. Process. **8** (2010), no. 3, 431–455.

[30] M. Ehler and K. Okoudjou, *Minimization of the probabilistic p-frame potential*, J. Stat. Plann. Inference **142** (2012), no. 3, 645–659.

[31] P. Elias, *Error-correcting codes for list decoding*, IEEE Trans. Inform. Theory **37** (1991), no. 1, 5–12.

[32] V. Elser and R. P. Millane, *Reconstruction of an object from its symmetry-averaged diffraction pattern*, Acta Crystallographica Section A **64** (2008), no. 2, 273–279.

[33] F. Filbir and H. N. Mhaskar, *A quadrature formula for diffusion polynomials corresponding to a generalized heat kernel*, J. Fourier Anal. Appl. **16** (2010), no. 5, 629–657.

[34] ——, *Marcinkiewicz–Zygmund measures on manifolds*, Journal of Complexity (2011).

[35] W. Fulton, *Intersection theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete, vol. 2, Springer-Verlag, 1984.

[36] Q. T. Le Gia and H. N. Mhaskar, *Localized linear polynomial operators and quadrature formulas on the sphere*, SIAM J. Numer. Anal. **47** (2008), no. 1, 440–466.

[37] R. Goodman and N. R. Wallach, *Representations and invariants of the classical groups*, Encyclopedia of Mathematics and its Applications, vol. 68, Cambridge University Press, 1998.

[38] D. Gross, *Recovering low-rank matrices from few coefficients in any basis*, IEEE Trans. Inform. Theory **57** (2011), 1548–1566.

[39] P. C. Hansen, *Analysis of discrete ill-posed problems by means of the L-curve*, Siam Review **34** (1992), no. 4, 561–580.

[40] R.B. Holmes and V. Paulsen, *Optimal frames for erasures*, Lin. Alg. Appl. **377** (2004), 31–51.

[41] J. Kovačević and A. Chebira, *Life beyond bases: The advent of frames (part i)*, IEEE Signal Processing Magazine **24** (2007), no. 4, 86–104.

[42] ______, *Life beyond bases: The advent of frames (part ii)*, IEEE Signal Processing Mag. **24** (2007), no. 5, 115–125.

[43] G. Kutyniok, A. Pezeshki, R. Calderbank, and T. Liu, *Robust dimension reduction, fusion frames, and Grassmannian packings*, Appl. Comput. Harmon. Anal. **26** (2009), no. 1, 64–76.

[44] L. B. Montefusco and S. Papi, *A parameter selection method for wavelet shrinkage denoising*, BIT **43** (2003), 611–626.

[45] J. M. Renes, R. Blume-Kohout, A. J. Scott, and C. M. Caves, *Symmetric informationally complete quantum measurements*, J. Math. Phys. **45** (2004), 2171–2180.

[46] A. J. Scott, *Tight informationally complete quantum measurements*, Journal of Physics A-Mathematical and General **39** (2006), no. 43, 13507–13530.

[47] F. Sottile, *From enumerative geometry to solving systems of polynomial equations*, vol. 2, Springer-Verlag, 2001.

[48] R. Vershynin, *Introduction to the non-asymptotic analysis of random matrices*, Compressed sensing, Theory and Applications (Y. Eldar and G. Kutyniok, eds.), Cambridge University Press, 2012, pp. 210–268.